

Preliminary version
May 1993

Scattering of very light charged particles

J C Taylor

DAMTP, Cambridge University, Silver Street, Cambridge, UK

I advance arguments against the view that the Lee-Nauenberg-Kinoshita theorem is relevant in practice to the scattering of charged particles as their mass tends to zero. I also discuss the case of massive coloured particle scattering.

Scattering cross-sections for reactions involving, in the initial state, a charged particle of mass m contain (in perturbation theory) $\ln m$ terms, which become large as $m \rightarrow 0$. These logarithms come from virtual photons which are nearly parallel to the charged particle. In 1964, following earlier work of Kinoshita [1], Lee and Nauenberg [1] proved that by summing over an appropriate set of initial states the $\ln m$ terms could be removed. Since then it seems not to have been generally agreed whether the initial states required for the theorem correspond to physically realistic situations. (For further references, see [2].)

In 1992, Contopanagos and Einhorn [2] (referred to as CE below) published a paper which, amongst many other things, seemed to claim with some certainty that the Lee-Nauenberg-Kinoshita (KLN) initial-state sum does represent physical reality. They went further and studied qualitatively the extent of the initial-state sum necessary to represent a typical physical, realistic situation. This has the great virtue of focusing the argument in a concrete way.

I remain unconvinced of the physical relevance of the initial-state sums, and, lest the very thorough CE paper should be thought to close the argument, I write this note to emphasize the questions which, to my mind, remain. I do not claim any complete understanding, but I hope this note may at any rate provoke further discussion.

In QED, soft divergences are well understood in the Bloch-Nordsieck theory. Further, the application of the KLN theorem to final-state collinear divergences is uncontroversial. So I concentrate on initial-state collinear divergences in QED. I also discuss the relevance of the KLN theorem to soft divergences in (perturbative) QCD, where, even with massive particles in the initial state, there are infrared divergences (uncancelled by Bloch-Nordsieck *final*-state sums) [3].

First note that the opinion of CE is not totally clear. In the second paragraph of the paper they state ‘we shall show that the requisite initial-state sum does inevitably occur in massless theories’. But in Section 4 they say: ‘The equality displayed in Eq. (4.1) requires a specific relative weighting among degenerate initial states, viz., the same

phase space normalizations that apply to final states. While this relation is an indisputable mathematical fact, it carries the paradoxical implication that initial-state degeneracy is to be associated with a certain relative weight between, say, an incoming single electron of definite energy and an electron of much lower energy accompanied by a hard but nearly collinear photon. This conflicts with the intuitive notion of an electron beam as well as the idea that one may prepare arbitrary linear combinations of states in Hilbert space. A complete resolution of this paradox requires a more careful analysis of the measurement process. While we have not carried out such a study, we believe it would show that ...' (This understates the paradox: the relative *phases* as well as weights of the degenerate states must be right.) The general tone of the CE paper, however, seems to be one of great confidence in physical relevance of the initial-state sums.

Something very strong is being claimed: that whatever the physical situation, collider or fixed target, etc., there will be some KLN initial-state (with some choice of parameters) which corresponds to it.

I begin by briefly summarizing CE's formulation, which has the advantage of being rather concrete. In the spirit of CE, I will consider an 'electron' of very small mass m , rather than a truly massless one. This avoids mathematical questions about Hilbert spaces.

CE construct two asymptotic Hamiltonians, $H_A(\delta)$ and $H_A(\delta')$, where δ and δ' are each some sort of 'resolution' parameter (or set of parameters) connected with the initial and final states. They then define two Møller operators

$$\Omega^+ = \lim_{t \rightarrow -\infty} e^{iHt} e^{-iH_A(\delta)t}, \quad (1)$$

$$\Omega^- = \lim_{t \rightarrow +\infty} e^{iHt} e^{-iH_A(\delta')t}. \quad (2)$$

(The H_A are supposed to be chosen such that these limits exist.) Lastly they define an operator

$$S_A(\delta, \delta') = \Omega^- S \Omega^{+\dagger}, \quad (3)$$

where S is the ordinary Feynman-Dyson scattering operator.

Then the following claims are made about S_A :-

- (a) The matrix elements of S_A between ordinary Fock states are insensitive to m for very small m (i.e. $m/E \ll \delta, \delta'$).
- (b) These matrix elements describe realistic scattering experiments, provided δ and δ' are chosen suitably.
- (c) There is no ambiguity about the Fock states to be chosen. For example, if we choose, instead of a single electron state, an electron-photon state within the resolution angle δ , the result is zero.
- (d) For the example of an electron beam in a collider, δ is to be identified, at least in order of magnitude, with r/L , where L is the distance from the final focus (FF) to the intersection point (IP), and r is a dimension of the beam spot at IP. At LEP, for example, CE estimate that $\delta \simeq 3 \times 10^{-6} < m/E \simeq 10^{-5}$; so that the claimed KLN cancellation is not far from being of practical importance.

Presumably the interpretation is that the coherent superposition of Fock states in, for example,

$$\Omega^{+\dagger}|\text{one-electron, } \mathbf{p}\rangle \quad (4)$$

is automatically and inevitably generated in the machine at FF or upstream of it.

The reason for believing this is, I suppose, that the mechanism for producing the beam could be analysed in the same terms, using the operator S_A in (3), and then the final states of that production process would automatically be described in a form involving the $\Omega^\pm$ operators, like (4).

My first objection to this is that the beam emerging from the production process, considering that as a scattering process of some kind, would be an *out* state; and so of the form

$$\Omega^{-\dagger}|\text{one-electron, } \mathbf{p}\rangle, \quad (5)$$

rather than (4). To express (5) in terms of states like (4) requires the ‘collinear S-operator’

$$\sigma = \Omega^{-\dagger}\Omega^+, \quad (6)$$

which describes the scattering of one collinear state into another, i.e. the emission and absorption of collinear photons by the electron. The operator σ is not unity, indeed its matrix elements in general themselves contain collinear divergences.

There is another, related, objection. The analysis has all been done in terms of plane-wave scattering states. This is of course an idealization. To be realistic one needs wave-packet states of some kind. For example, the initial beam should really be described by a wave-packet, with a transverse size of order, say, r' . Consider now a one-electron-one-photon Fock state

$$|\mathbf{p} - \mathbf{k}, \mathbf{k}\rangle \quad (7)$$

contained in (5). Again, this should really consist of wave-packets with limited transverse extent. Certainly this would be true for photons radiated due to the focusing magnetic field. The photon wave packet is diverging from the electron one at an angle of the order of the angle θ between $\mathbf{p}$ and $\mathbf{k}$, where $\theta < \delta$. At the IP, therefore, the centres of the two wave-packets will be separated by a transverse distance of order $L\alpha$ which has a range up to a maximum of order r . However, the state required in (4), being the time-reverse of an out-state, requires the two wave-packets to be converging and to exactly overlap at IP. Anything less than exact overlap would result in incomplete cancellation of the collinear divergences in (3).

I have tried to articulate my unease about the assumptions made in [2], especially the argument for the relevant magnitude of δ . I do not deny that sort of effect is to be expected when Lm/E becomes *microscopic*. The electron propagator probably does not have a pole at $E = (\mathbf{p}^2 + m^2)^{\frac{1}{2}}$, but a branch point there. An electron propagating over a distance L probably samples a length of the branch-cut of order $1/L$. This would correspond to an opening angle $\delta \simeq (EL)^{-1/2}$.

In an accompanying paper [4], CE analyse ‘evanescent’ processes, such as helicity flip of an electron emitting a collinear photon. Although the matrix element is proportional to m , the phase space has a factor $1/m^2$, and so the rate appears to be finite as $m \rightarrow 0$.

CE claim that KLN initial states will cancel this effect when $m/E < \delta$. My remarks above apply equally to the order of magnitude of δ here.

Finally I briefly mention a nonabelian case. Here there are soft divergences, uncanceled by the final-state Bloch-Nordsieck mechanism, when there are coloured particles in the initial state even if these have mass [3]. For example, one could take $b\bar{b}$ reactions in a hypothetical unconfined world. To remove these soft divergences, the coherent initial states would have to include soft gluons moving in all directions: collinearity with the quarks is not relevant. It is difficult to see how the ‘accelerator’ producing the quark beams could also produce coherent gluons converging on the annihilation point from *all* directions.

This example may not be so removed from physics. An extension of the QCD factorization theorem to higher-twist would require a meaning to be given to quark cross-sections. (The problem only appears at higher-twist, because the uncanceled soft divergences are suppressed by a factor E^{-2} .)

I am grateful to D.R.T. Jones for discussions on the subject, but he does not have any responsibility for the shortcomings of this note.

References

- [1] T. Kinoshita, J. Math. Phys. **3**, 650 (1962); T.D. Lee and M. Nauenberg, Phys. Rev. **133**, B1549 (1964).
- [2] H.F. Contopanagos and M.B. Einhorn, Phys. Rev. **D45**, 1291 (1992).
- [3] R. Doria, J. Frenkel and J.C. Taylor, Nucl. Phys. **B168**, 93 (1980); A. Andrasi *et al*, Nucl. Phys. **B183**, 445 (1981); C. Di’Lieto *et al*, Nucl. Phys. **B183**, 223 (1981).
- [4] H.F. Contopanagos and M.B. Einhorn, Phys. Rev. **D45**, 1322 (1992).